

② $s_1, s_2, \dots, s_N \in H^0(X, mL)$

define $\varphi_\alpha = \log \left(\sum_{i=1}^N |s_{i,\alpha}|^{2/m} \right)^{1/2}$, then $\varphi_\alpha = \varphi_\beta + \log |\varphi_\beta|$, $\varphi_\alpha \in \text{psh}(V_\alpha)$.

③ Let $s_1^{(m)}, \dots, s_{N_m}^{(m)} \in H^0(X, mL)$, $m \in \mathbb{N}$

define $\varphi_{\min, \alpha}^{\text{Siu}} = \log \left(\sum_{m=1}^{\infty} \theta_m \sum_{k=1}^{N_m} |s_{k,\alpha}|^{2/m} \right)^{1/2}$, where $\{\theta_m > 0\}$ is a suitable sequence s.t. $\sum_{m=1}^{\infty} \theta_m \sum \dots$ converges.

Prbl./Exer.*:

Let $X/\mathbb{C}$ be a proj. mfd, L a psef. line bdl on X , $\theta \in G(L)$ a s.m. rep.

Let $\varphi_{\min}^{\text{Demailly}} = \sup \{ u \mid \theta + \frac{i}{\pi} \partial \bar{\partial} u \geq 0 \text{ in the sense of currents} \}$

Let $s_1^{(m)}, \dots, s_{N_m}^{(m)} \in H^0(X, mL)$ be the basis of $H^0(X, mL)$, $m \in \mathbb{N}$.

Then $R(L) = \bigoplus_{m=1}^{\infty} H^0(X, mL)$ is finitely generated $\iff \varphi_{\min}^{\text{Demailly}} = \varphi_{\min}^{\text{Siu}} + o(1)$.

#

Multiplier ideal sheaves:

Local case. $\Omega \subset \mathbb{C}^n$, $\varphi \in \text{psh}(\Omega)$

Def. the multiplier ideal sheaf associated to φ is defined by

$\mathcal{I}(\varphi) = \bigcup_{x \in \Omega} \mathcal{I}(\varphi)_x$, where $\mathcal{I}(\varphi)_x = \{ f \in \mathcal{O}_x \mid |f|^2 e^{-2\varphi} \in L^1_x \}$.

Thm. $\mathcal{I}(\varphi)$ is coherent on Ω . Moreover, if Ω is a bounded Stein open set, then $\mathcal{I}(\varphi)$ is generated by any Hilbert basis of the L^2 -space $H^2(\Omega, \varphi) = \{f \in \mathcal{O}(\Omega) \mid \int_{\Omega} |f|^2 e^{-2\varphi} dv < \infty\}$.

Proof: The statement is local, w.l.o.g., assume Ω is an open ball in $\mathbb{C}^n$.
(or any bdd. pseudoconvex open set)

$H^2(\Omega, \varphi)$ is a Hilbert space. Let $\{e_k\}_{k=1}^{\infty}$ be a basis; $\mathcal{F}^l$ = the sheaf generated by $\{e_1, e_2, \dots, e_l\}$.

Then $\mathcal{F}^l$ is coherent for any $l \in \mathbb{N}$.

$$\mathcal{F}^1 \subseteq \mathcal{F}^2 \subseteq \dots \subseteq \mathcal{F}^l \subseteq \dots \subseteq \mathcal{O}_{\Omega}$$

Strong Noetherian property of coherent sheaves $\Rightarrow$ on any $K \subset \subset \Omega$, the sequence is stationary for $l \geq l_0$.

$\Rightarrow$ the sheaf $\mathcal{I}$ generated by $H^2(\Omega, \varphi)$ is coherent.

Claim: $\mathcal{I}(\varphi) = \mathcal{I}$.

It's clear that $\mathcal{I} \subseteq \mathcal{I}(\varphi)$, we need to verify that $\mathcal{I}(\varphi) \subseteq \mathcal{I}$.

This follows from Nakayama's lemma:

[A : a local ring, E : a f.g. A -module, $F \subseteq E$: a submodule, $\mathfrak{m}$: maximal ideal]
then $E = F + \mathfrak{m} \cdot E \Rightarrow E = F$

In our setting let $E = \mathcal{I}(\varphi)_x$, $F = \mathcal{I}_x$, $A = \mathcal{O}_x$, $\mathfrak{m} = \mathfrak{m}_x$.

We need to verify that $\mathcal{I}(\varphi)_x = \mathcal{I}_x + \mathfrak{m}_x \cdot \mathcal{I}(\varphi)_x$

Indeed, we will prove that: $\forall s \in \mathbb{N}_{\geq 0}$, $\mathcal{I}(\varphi)_x = \mathcal{I}_x + \mathfrak{m}_x^{s+1} \mathcal{I}(\varphi)_x$.

更正: 下面实际证明的是 $\mathcal{I}(\varphi)_x = \mathcal{I}_x + \mathfrak{m}_x^{s+1} \cap \mathcal{I}(\varphi)_x$,
应用的交换代数结果为 Krull lemma.

Let $f \in \mathcal{J}(\varphi)_x$ in some nbhd. V of x , Θ : a cutoff function with $\text{supp } \Theta \subset V$ 18
 s.t. $\Theta = 1$ near x

We solve the equation $\bar{\partial} u = \bar{\partial}(\Theta f)$ with metric weight $\tilde{\varphi} = \varphi + (1+s) \log |z-x| + |z|^2$
 (Hörmander's L^2 -existence thm)

$\Rightarrow$ the equation has a solution u satisfying:

$$\int_{\Omega} |u|^2 e^{-2\tilde{\varphi}} \leq \int_{\Omega} |\bar{\partial}(\Theta f)|^2 e^{-2\tilde{\varphi}} < \infty$$

$$\int_{\Omega} |u|^2 e^{-2\varphi} \cdot |z-x|^{-2(1+s)}$$

$$\Rightarrow u_x \in \mathcal{J}(\varphi)_x \cdot \mathcal{M}_x^{s+1}$$

Let $F = \Theta f - u$, then $F \in H^2(\Omega, \varphi)$ and $F_x - f_x = -u_x$
 $\cap \quad \cap$
 $\mathcal{J}(\varphi)_x \quad \mathcal{J}(\varphi)_x \mathcal{M}_x^{s+1}$

Lem. $F_x \in \mathcal{J}_x$.

This follows from the thm on closure of module:

Any submodule $M \subseteq \mathcal{O}_x^{\oplus p}$ is closed in the following sense: if $F \in \mathcal{O}_x^{\oplus p}(V)$

and $\exists f_k \in \mathcal{O}^{\oplus p}(V)$ s.t. $f_k \xrightarrow{C_{loc}} F$ and the germ of $f_k \in M$

then the germ of $F \in M$.

Applying this to $M = \mathcal{J}$ shows that $F \in \mathcal{J}_x$. #

Exercise 1. Provide the details of the proof on the above lemma and closure of modules theorem.
 (ref. [Gunning-Rossi, Analytic Functions of Several Complex Variables, chap 2, sect. E])

global case of multiplier ideal sheaves.

19

$(X/\mathbb{C}, L, h)$: By the def. of psh line bundle (L, h) ($\Leftrightarrow$ metric weights $\varphi_\alpha \in \text{psh}(V_\alpha)$)
 $\varphi_\alpha = \varphi_\beta + \log |g_{\alpha\beta}|$ on $V_\alpha \cap V_\beta$


For $x \in V_\alpha \cap V_\beta$, it is clear that $J(\varphi_\alpha)_x = J(\varphi_\beta)_x = J(h)_x$.

Def. The multiplier ideal sheaf $J(h)$ associated to (L, h) is given by $J(h) = \bigcup_{x \in X} J(h)_x$.

Note that singular metrics can be defined on $\mathbb{R}$ -line bundles by replacing $|g_{\alpha\beta}|$ by some real combination [#]

Examples

① $X/\mathbb{C}$, $D = \sum_{j=1}^k \alpha_j D_j$: effective $\mathbb{R}$ -divisor with SNC sp. D :



Associated to D this is a singular metric φ_D : locally $D_j = g_j^{-1}(0)$ for some hol. function g_j

$\varphi_D = \sum_{j=1}^k \alpha_j \log |g_j|$ — these φ_D patch to a metric weight on the $\mathbb{R}$ -line bundle $\mathcal{L}_D$

$$\mathcal{L}_D = \sum_{j=1}^k \alpha_j \mathcal{L}_{D_j}.$$

Let's compute $J(\varphi_D)$:

$$f \in J(\varphi_D)_x \Leftrightarrow \int_{V_x} |f|^2 e^{-2\varphi_D} = \int_{V_x} |f|^2 / \prod_{j=1}^k |g_j|^{2\alpha_j} < \infty \quad (*)$$

D SNC $\Rightarrow$ for every x , the defining functions g_j complete to a local coordinate system $(z_1, z_2, \dots, z_n)$

Exercise 2.

Exer : $(*)$ holds $\Leftrightarrow f$ is divisible by $\prod_j g_j^{m_j}$ where $m_j - \alpha_j > -1$
 (i.e., $m_j \geq \lfloor \alpha_j \rfloor$)

Therefore, $J(\varphi_D) = \mathcal{O}(-\sum_{j=1}^k |\alpha_j| D_j) = \mathcal{O}(-LD)$.

注记：以下关于push-forward后面的层，记号有些混乱，请自行更正。

#

How about the case when D is NOT SNC? This follows from the projection formula:

Let $\mu: \hat{X} \rightarrow X$ be a modification between complex mfd's, $\varphi \in \text{psh}(X)$, then

$J(\varphi) = \mu_* \mathcal{O}_{\hat{X}}(K_{\hat{X}/X} \otimes J(\varphi \circ \mu))$, where $K_{\hat{X}/X} = K_{\hat{X}} - \mu^* K_X$ is the relative canonical sheaf.

$(\Leftrightarrow K_X \otimes J(\varphi) = \mu_*(K_{\hat{X}} \otimes J(\varphi \circ \mu)).$

Proof. Exer. Hint: $f \in J(\varphi)_X \Leftrightarrow \int_U |f|^2 e^{-2\varphi} dz \wedge \bar{z} < \infty$, by using the identification

$f(z) \mapsto f(z) dz$, $f \in J(\varphi)_X$ can be considered as a local section of $K_X \otimes J(\varphi)$

show that $K_X \otimes J(\varphi) \xrightarrow{\sim} \mu_*(K_{\hat{X}} \otimes J(\varphi \circ \mu))$ $(\mu: \hat{X} \rightarrow X)$
 $\mu^* \omega \mapsto \omega$
 $f(z) dz \mapsto \mu^* f(\omega) \text{Jac}(\mu) \cdot d\omega$, $\text{Jac}(\mu) = \frac{\partial \mu}{\partial \bar{\omega}} = \left[\frac{\partial^2 \mu}{\partial \bar{\omega} \partial \omega} \right]$

#

Exercise 3. Provide the details for proof of the projection formula.

② $X/\mathbb{C}$, $D = \sum_{j=1}^k \alpha_j D_j$: effective IR-divisor. $(\varphi_D = \sum \alpha_j \log |g_j|)$

Black box : Hironaka's desingularization theorem.

Consider a log-resolution of (X, D) , i.e. a modification $\mu: \hat{X} \rightarrow X$ satisfying

• $\hat{X}$ smooth

• $\mu^* D + \text{Exc}(\mu)$ SNC (associated to $\mu^* D$ is $\varphi_{D \circ \mu}$)

By the projection formula,

$$\begin{aligned} J(\varphi_D) &= \mu_* (\mathcal{O}_{\hat{X}/X} \otimes J(\varphi_D \circ \mu)) \\ &= \mu_* (\mathcal{O}_{\hat{X}/X} - \mu^* D). \end{aligned}$$

Rem. on the right hand side, μ can be different, but $J(\varphi_D)$ is fixed.

• analytic multiplier ideal sheaf $\leftrightarrow$ algebraic multiplier ideal sheaf.

Def. (PIAG, vol 2, chap. 9)


s.m. complex var.

$X/\mathbb{C}$: ~~proj. mfd.~~ D : effective $\mathbb{Q}$ -divisor, $\mu: \hat{X} \rightarrow X$: a log resolution of (X, D)

then $J(D)$ is defined by $J(D) = \mu_* (\mathcal{O}_{\hat{X}} (K_{\hat{X}/X} - \mu^* D))$

(In particular, $J(\varphi_D) = J(D)$.)

Ex. $X = \mathbb{P}^2$, $D = \frac{2}{3}(A_1 + A_2 + A_3)$, where A_1, A_2, A_3 are 3 distinct lines through 0.



Thus, $\mu^* D = \frac{2}{3} (A_1' + A_2' + A_3') + 2E$

Note that $K_{\hat{X}/X} = E$, we get $J(X, D) = J(\varphi_D) = \mu_* (\mathcal{O}_{\hat{X}} (-E))$

↑
maximal ideal of hol. functions
vanishing at 0.

More Examples:

22

$\mathfrak{a} \subseteq \mathcal{O}_X$: an ideal sheaf, locally $\mathfrak{a} = \langle f_1, f_2, \dots, f_N \rangle$.

For $c > 0$, we define $\varphi_{\mathfrak{a}^c} = \frac{1}{2} \log (|f_1|^2 + \dots + |f_N|^2)^c$.

Prop/Exer. If $g_1, \dots, g_M$ is another family of local generators of $\mathfrak{a}$, then we get another

psd function $\tilde{\varphi}_{\mathfrak{a}^c} = \frac{1}{2} \log \left(\sum_{k=1}^M |g_k|^2 \right)^c$, we have $\tilde{\varphi}_{\mathfrak{a}^c} = \varphi_{\mathfrak{a}^c} + O(1)$.

Therefore $J(\varphi_{\mathfrak{a}^c}) = J(\tilde{\varphi}_{\mathfrak{a}^c})$.

algebra-geometrical def: Let $\mu: \hat{X} \rightarrow X$ be a log resolution of $(X, \mathfrak{a})$, i.e. a birational morphism satisfying:

- $\hat{X}$ sm.
- $\mu^* \mathfrak{a} = \mathfrak{a} \cdot \mathcal{O}_{\hat{X}} = \mathcal{O}(-F)$ with $F + \text{Exc}(\mu)$ SNC

$$J(X, \mathfrak{a}^c) := \mu_* \mathcal{O}_{\hat{X}}(K_{\hat{X}/X} - \lfloor cF \rfloor).$$

CLAIM: $J(X, \mathfrak{a}^c) = J(\varphi_{\mathfrak{a}^c})$

Exercise 4. Prove the claim. (hint: apply the projection formula to analytic multiplier ideal sheaf and the property of log-resolution.)

Lehman numbers and $J(\varphi)$.

23

Recall, given $\varphi \in \text{PSH}(\Omega)$, $x \in \Omega$,
$$V(\varphi, x) = \sup \{ t \geq 0 \mid \varphi(z) \leq t \log |z-x| + O(1) \text{ around } x \}$$

$\Rightarrow$ Prop./Exer.: If $V(\varphi, x) \geq n+s$ for some $s \in \mathbb{Z}_{\geq 0}$, then $J(\varphi)_x \subseteq m_x^{s+1}$.

On the other hand,

Thm (Skoda 1972) if $V(\varphi, x) < 1$, then $e^{-2\varphi} \in L_x^1$, in particular, $J(\varphi)_x = \mathcal{O}_x$. #

Denote $E_c(\varphi) = \{ x \in \Omega \mid V(\varphi, x) \geq c \}$, $c > 0$

$V(J(\varphi)) = \text{the zero variety of } J(\varphi) = \{ x \in \Omega \mid e^{-2\varphi} \notin L_x^1 \}$

then we get: $E_n(\varphi) \subseteq V(J(\varphi)) \subseteq E_1(\varphi)$.

Thm. (Boucksom-Favre-Jonsson) + (Guen-Zhou's openness thm)

Let $\varphi_1, \varphi_2 \in \text{PSH}(\Omega)$, $x \in \Omega$, ~~then: $\pi: \hat{\Omega} \rightarrow \Omega$ blow-up at x~~

TFAE:

• $\forall t > 0$, $J(t\varphi_1)_x = J(t\varphi_2)_x$

• $\forall$ proper modification $\pi: \Omega_\pi \rightarrow \Omega$ over x and any p.t. $p \in \pi^{-1}(x)$

$V(\varphi_1 \circ \pi, p) = V(\varphi_2 \circ \pi, p)$.

#